


A P P E N D I X IV:

THE CURRENT CLAIMS (clean version):

1. (previously presented) A compound of formula I



(I)

in which

R¹ represents a hydrogen or a methyl group;

R² represents a hydrogen atom or an optionally substituted C₁-C₁₀-alkyl, C₂-C₁₀-alkenyl, C₂-C₁₀-alkynyl, C₄-C₁₀-alkadienyl or phenyl group, wherein the optional substituents are selected from the group consisting of nitro, cyano, C₃-C₆-cycloalkyl, C₃-C₆-cycloalkenyl, C₁-C₆-haloalkyl, C₃-C₆-halocycloalkyl, C₁-C₆-alkoxy, C₁-C₆-haloalkoxy, tri-C₁-C₄-alkylsilyl, phenyl, halophenyl, dihalophenyl and pyridyl;


Hal represents a halogen atom; and

L¹ through L⁵ each represent a hydrogen or halogen atom or an C₁-C₁₀-alkyl, C₁-C₁₀-alkoxy or nitro group, provided that at least one of L¹ through L⁵ represents a nitro or alkoxy group.

2. (original) A compound according to claim 1 in which at least one of L¹ and L⁵ represents a halogen atom.
3. (original) A compound according to claim 1 in which R² represents a hydrogen or a C₁₋₁₀ alkyl group.
4. (original) A compound according to claim 1 in which at least one of R¹ and R² represents a hydrogen atom.
5. (previously presented) The compound of formula I defined in claim 1 which is selected from the group consisting of
 5-chloro-6-(4-methoxyphenyl)-7-(2,2,2-trifluoroethylamino)-[1,2,-4]triazolo[1,5-a]pyrimidine;
 5-chloro-6-(4-nitrophenyl)-7-(2,2,2-trifluoroethylamino)-[1,2,4]-triazolo[1,5-a]pyrimidine; and

5-chlor -6-(2,6-difluoro-4-methoxyphenyl)-7-[2-(1,1,1-trifluoro)-propyl)amino]-[1,2,4]triazolo[1,5-a]pyrimidine.

6. (previously presented) A process for the preparation of a compound of formula I as defined in claim 1, which process comprises:
treating a compound of formula II



with an amine of formula III



in which M represents a hydrogen atom or a metal atom,
to produce the compound of formula I.

7. (previously presented) A fungicidal composition which comprises a carrier, and as active agent, at least one compound of formula I as defined in claim 1.
8. (previously presented) A method of combating fungus at a locus which comprises treating the locus with a fungicidally effective amount of a compound of formula I as defined in claim 1.
9. (currently amended) The process of claim 6 wherein the metal atom represented by M is selected from the group consisting of Li, Na, K, Zn and Cu.